Open formula purified diets for lab animals



Product Data - D12266B



gm%

18.5

56.7

15.6

4.41

Total

kcal/gm

kcal%

16.8

51.4

31.8

100

4595.8

Description Purified match to C11024, the "condensed milk" diet.

number and expiration date.

Gamma-Irradiation

Pellet, Powder, Liquid

Yes. Add 10 days to delivery time.

Most diets require storage in a cool

dry environment. Stored correctly

they should last 6 months.

Control Diets

D12489B

Used in Research

Obesity Diabetes Hypertension

Lead Time

Form

Shelf Life

5-7 business days.

Carbohydrate Fat Packaging Product is packed in 12.5 kg box. Each box is identified with the product name, description, lot

Formula

Protein

Total

Product # D12266B

Ingredient	gm	kcal
	100	7(0
Casein, 30 Mesh	190	/60
DL-Methionine	3	12
Corn Starch	215	860
Maltodextrin 10	75	300
Sucrose	290	1160
		0
Cellulose, BW200	30	0
Butter Fat. Anhydrous	44.2	397.8
Corn Oil	118	1062
	110	1002
Mineral Mix S10001	40	0
Calcium Carbonate	5.5	0
Sodium Chloride	5.5	0
Potassium Citrate, 1 H2O	13.5	0
Vitamin Mix V10001	11	44
Choline Bitartrate	2	0
FD&C Yellow Dye #5	0	0
FD&C Red Dve #40	0.1	0
	0.1	0



Research Diets, Inc. 20 Jules Lane New Brunswick, NJ 08901 USA Tel: 732.247.2390 Fax: 732.247.2340 info@researchdiets.com

1042.8



- 1. Alberts, P. et al. Selective inhibition of 11ß-hydroxsteroid dehydrogenase type 1 decreases blood glucose concentrations in hyperglycaemic mice.
- Diabetologia. 45:1528-1532, 2002.
 Alberts, P. et al. Selective inhibition of 11ß-hydroxysteroid dehydrogenase type 1 improves hepatic insulin sensitivity in hyperglycemic mice strains. Endocrinology. 144:4755-4762, 2003.
- 3. Turnbull, A.V., et al. Selective antagonism of the NPY Y5 receptor does not have a major effect on feeding in rats. Diabetes. 51:2441-2449, 2002.
- 4. Block, M.H., et al. Discovery and optimization of a series of carbozole ureas as NPY5 antagonists for the treatment of obesity. J. Med. Chem. 45:3509-3523, 2002.
- 5. Boustany, C.M., et al. Activation of the systemic and adipose rennin-angiotesin system in rats with diet-induced obesity and hypertension. Am. J. Physiol. Integr. Comp. Physiol. 287: R943-R949, 2004.
- Chandler, P.C., et al. Change in CCK-8 response after diet-induced obesity and MC3/4-receptor blockade. Peptides. 25:299-306, 2004.
- Chandler, P.C., et al. Feeding response to melanocortin agonist predicts preference for and obesity from a high-fat diet. Physiology and Behavior. 85:221-230, 2005.
- 8. Chen, A.S., et al. Role of the melanocortin-4 receptor in metabolic rate and food intake in mice. Transgenic Research. 9:145-154, 2000.
- 9. Clegg, D.J., et al. Reduced anorexic effects of insulin in obesity-prone rats fed a moderate-fat diet. Am. J. Physiol. 288:981-986, 2005.
- 10. DeRuisseuu, L.R., Parsons, A.D., & Overton, J.M. Adaptive thermogenesis is intact in B6 and A/J mice studied at thermoneutrality. Metabolism. 53:1417-1423, 2004.
- Bobrian, A.D., et al. Effect of salt on hypertension and oxidative stress in a rat model of diet-induced obesity. Am. J. Physiol. Renal Physiol. 285::F619-F628, 2003.
- 12. Dobrian, A.D., et al. Development of Hypertension in a rat model of diet-induced obesity. Hypertension. 35:1009-1015, 2000.
- 13. Dobrian, A.D., et al. Oxidative stress in a rat model of obesity-induced hypertension. Hypertension. 37:554-560, 2001.
- 14. Dobrian A.D. et al. Pioglitazone prevents hypertension and reduces oxidative stress in diet-induced obesity. Hypertension. 43: 48-56, 2004.
- 15. Ghibaudi, L., et al. Fat intake affects adiposity, comorbidity factors, and energy metabolism of Sprague-Dawley rats. Obesity Research. 10:956-963, 2002.
- 16. Heinrichs, Stephen. Nonexercise muscle tension and behavioral fidgeting are positively correlated with food availability/palatability and body weight in rats. Physiology & Behavior. 79:199-207, 2003.
- 17. Larden, P.J., et al. Differential influence of peroxisome proliferators-activated receptors ? and -a on food intake and energy homeostasis. Diabetes. 52:2249-2259, 2003.
- Lauterio, T. J., et al. Plasma growth hormone secretion is impaired in obesity-prone rats before onset of diet-induced obesity. American Journal of Physiology. 275:E6-E11, 1998.
- Lauterio, T.J., Bond, J.P. * Ulman, E.A. Development and characterization of purified diet to identify obesity-susceptible and resistant rat populations. Journal of Nutrition. 124:2172-2178, 1994.
- 20. Lauterio, T.J., et al. Neuropeptide Y expression and endogenous leptin concentrations in a dietary model of obesity. Obesity research. 7:498-505, 1999.
- 21. Marsh, D.J., et al. Melanin-concentrating hormone 1 receptor-deficient mice are lean, hyperactive, and hyperphagic and have altered metabolism.
- PNAS. 99:3240-3245, 2002.
- 22. Shen, C.P., et al. Plasma agouti-related protein level: a possible correlation with fasted and fed states in humans and rats. Journal of Neuroendocrinology. 14:607-610, 2002.
- 23. Singh, K.A., et al. Acute insulin-induced elevations of circulating leptin and feeding inhibition in lean but not obese rats. Am. J. Regul. Integr. Comp. Physiol. 289:373-379,2005.
- 24. Singh, D.K., et al. Phosphorylation of supernatant protein factor enhances wits ability to stimulate microsomal squalene monooxygenase. The Journal of Biological achemistry. 278:5646-5651, 2003.
- 25. Strack, A.M. et, al. Regulation of body weight and carcass composition by sibutramine in rats. Obesity Research. 10:173-181, 2002.
- 26. Turnbull, A.V. Selective antagonism of the NPY Y5 receptor does not have a major effect on feeding in rats. Diabetes. 51:2441-2449, 2002.
- 27. Warden, C. H., et al. Identification of a congenic mouse line with obesity and body length phenotypes. DOI. 15:460-471, 2004.
- 28. Zhou, D., et al. Enhanced running wheel activity of both Mch 1r- and Pmch-deficient mice. Regulatory Peptides. 124:53-63, 2005.



Research Diets, Inc. 20 Jules Lane New Brunswick, NJ 08901 USA Tel: 732.247,2390 Fax: 732.247,2340 info@researchdiets.com