



Phytoestrogens and Chow

Report ▶ Repeat ▶ Revise

Phytoestrogens – Uninvited, Troublesome Guests in Scientific Research

by Lorene M. Leiter, Ph.D.

Project Manager and Scientist- Research Diets, Inc.

Eliminate variables. It's a fundamental rule in scientific research. Purified diets were designed for this very purpose – to minimize variables so that scientists can be confident they are studying only the nutrient or compound of interest. With the availability of this remarkable tool, it's hard to understand why so many researchers continue to use grain-based chow diets in nutrition studies and even, as we shall see, in studies that are not necessarily nutrition-oriented, such as drug and compound efficacy studies, or behavioral studies.

Perhaps many researchers are unaware of the profound differences between the two diets. Purified diets are made with refined ingredients, such as casein, corn starch, sucrose, corn oil, and a fixed amount of vitamins and minerals. A scientist has complete control over the ingredients added or subtracted (1).

In contrast, grain-based chow diets are made with ground plants, such as corn, wheat, oats, soybean, and alfalfa. There's a problem – plants vary nutritionally from harvest to harvest, and some, including soy and alfalfa, contain significant amounts of biologically active compounds called phytoestrogens, which also vary between harvests (2).


Phytoestrogens in chow add potentially powerful variables to experiments

Phytoestrogens are abundant in soy meal – the major protein source in most commonly-used laboratory chows; and a subclass of chemicals in the phytoestrogen family known as isoflavones are major culprits (3). Isoflavones interact with estrogen receptors and elicit estrogenic or anti-estrogenic effects (4). Therefore, they can interfere with studies influenced by exogenous estrogens. Estrogens regulate a variety of physiological pathways. They are involved in the growth and function of reproductive tissues of both males and females, as well as maintenance of the skeletal, central nervous and cardiovascular systems. Hence, isoflavones can interfere with countless studies and even render the interpretation of results meaningless.

To date, numerous studies in humans and animals leave little doubt that isoflavones in soy affect a variety of experimental endpoints. A brief mention of some examples follows, and extensive reviews on this topic have been published (5, 6, 7) FIG. 1.

Cancer. The low incidence of many cancers in Asian countries, where soy consumption is high, sparked an interest in soy's potential protective effects (8). As partial estrogen antagonists, isoflavones in soy were thought to play a role in preventing cancer. Indeed, laboratory research has confirmed the epidemiological findings. For example,

Phytoestrogens in my lab animal diet?



Do you know that grain-based diets contain significant and variable levels of phytoestrogens? If your studies involve pathways which are influenced by estrogenic activity, the diet you feed can have profound effects on your animal model at the molecular and cellular level.

Get your research under control with Research Diets phytoestrogen-free purified ingredient OpenSource Diets™.

Your results could depend on it.

rodents consuming isoflavones have fewer tumors and/or a delay in tumor development in mammary, liver, colon, and prostate cancer (9, 10, 11, 12). Experimental results sometimes conflict and the outcome likely depends on a variety of factors, including the isoflavone amount, duration of feeding, and hormonal state of the animal. Still, the evidence for isoflavones' role in cancer is mounting and must not be dismissed as insignificant in cancer research.

Metabolic Syndrome. Dietary soy has been shown to attenuate the development of hypertension in spontaneously hypertensive rats (13). Soy isoflavones have also been shown to reduce serum cholesterol and triglycerides (14, 15), and prevent the development of fatty liver (16). Rats fed high fat diets with soy as the protein source gained less weight and had less body fat compared with those fed high-fat diets with casein (17). Insulin-sensitivity was also improved in soy-fed mice (18).

Osteoporosis. Studies in both humans (post-menopausal women) and animal models of osteoporosis (ovariectomized rodents) show bone sparing effects of soy isoflavones (19, 20), and the results in animals are remarkably consistent.




FIGURE 1. Isoflavone content in a number of commonly used laboratory rodent diets. Each data point represents a batch of the diet formulations listed on the right. Diets P-Y are formulated without soy and have negligible amounts of isoflavones. Graphic representation - for details see reference (6).

Reproductive Development. Certain levels of isoflavones are capable of accelerating puberty and stimulating uterine growth in female rodents (21). In male rodents, a decrease in the weight of the prostate gland and sexual organs, as well as a decrease in testosterone, has been reported (22, 23). A literature review reveals inconsistent findings, and, as with other endpoints, likely depends on the dose and other factors. Yet, scientists seem to agree that dietary phytoestrogen content must be considered in studies directly or indirectly involving sexual development.

Behavior. Since estrogen receptors are expressed in the brain, it's no surprise that dietary isoflavones affect behavioral parameters. Effects on learning and memory, social interaction, anxiety-related behaviors, locomotor activity, and pain sensitivity have been reported (6, 24).

A cautionary tale

Mounting empirical evidence to date has implicated isoflavones in a wide range of experimental endpoints, leaving little doubt that, when choosing a diet, careful consideration must be given to the role of these compounds in mammalian physiology. After all, can meaningful conclusions really be drawn if the control diet itself has protective or adverse effects on the phenotype one is studying?

Nutrition scientists who, surprisingly, use grain-based diets as a control for purified diets must be made aware of these differences. In addition, the knowledge must be imparted to scientists studying

the efficacy of drugs on a variety of diseases that may be affected by compounds in soy. These include osteoporosis, cancer, heart disease, hypertension, pain sensitivity, and psychiatric disorders. Understandably, many researchers are reluctant to switch from grain-based chows to purified diets. They dare not change the diet in ongoing experiments, since these rely heavily on comparisons to historical data. Yet, a strong argument can be made for a break from the past: Was the "same diet" actually used over the years? The answer is no, since chows vary from batch to batch. Indeed, dietary phytoestrogen variability has likely caused a frustrating variability in data, or a misinterpretation of results, in many studies.

Does removing the phytoestrogen source from grain-based chow fix the problem?

In recognition of the large body of evidence of phytoestrogens' effects, some chow manufacturers now sell chow without soy meal or alfalfa. Instead, the ingredients include ground corn, wheat middlings, and ground wheat. The phytoestrogens in these diets are minimized but not eliminated. Importantly, these ingredients are not refined. Therefore, it is impossible to know which yet-to-be-discovered biologically active compounds are in the diet. Indeed, in 1940, an effect of phytoestrogens was noticed for the first time – when it was observed that sheep grazing on red clover, a phytoestrogen-rich plant, were less fertile (25).

It can certainly be argued that there's no such thing as a perfect diet in laboratory research, but at the very least, for experiments to be meaningful, we should strive to eliminate dietary variables, especially those known to elicit biological responses – namely, phytoestrogens.





Contact our Resource Center for valuable insight from more than 25 years of product experience in purified lab animal diets. Let us formulate the diets to meet your specific study needs.

Custom OpenSource Diets

Research Diets, Inc has pioneered the formulation and production of phytoestrogen- free diets in laboratory animals. Our scientists specialize in providing custom purified OpenSource diets. By carefully designing the diet formula to fit your protocol, you have complete control over small or large changes in diet composition. Plus you are able to report what your animals were fed, repeat the formula and revise diet composition as necessary.

Incorporate Test Compounds

Research Diets, Inc. will incorporate your test compound into pelleted diets for simple, safe dosing. Feeding test compounds eliminates dosing related stress to the animal, eliminates vehicle effects, and saves time and labor. Consult with one of our scientists on the formula, determine the dosage required and the diet will be produced and shipped in 5 to 7 business days.

Value Added Resource

The value of our products is in the scientific support we provide. Our Resource Center is staffed with Masters and Ph.D. level scientists with access to over 14,000 original formulas and a database of more than 3,800 journal articles. We welcome the opportunity to talk science with researchers throughout the world as we maintain our leadership role as the knowledge base for OpenSource Diet formulation.

BioDAQ Episodic Intake Monitor

BioDAQ E2 Episodic monitoring system measures the ad libitum food and water intake behavior of singly housed lab rats and mice at very high resolution in their home cage. Computer controlled electronics record food and water intake episodically by measuring the moment-to-moment, undisturbed intake behavior of the animals being studied. The animal's behavior defines the variable period for these intake measurements through their native behavior.



References

- Ricci MR and Ulman EA. Laboratory Animal Diets: a critical part of your in vivo research. *Animal Lab News* 4(6): 1-5, 2005.
- Thigpen JE, Setchell KDR, Ahlmark KB, Locklear J, Spahr T, Caviness GF, Goelz MF, Haseman JK, Newbold RR, and Forsythe DB. Phytoestrogen content of purified, open- and closed-formula laboratory animal diets. *Laboratory Animal Sciences* 49(5): 530-536, 1999.
- Brown NM and Setchell KDR. Animal models impacted by phytoestrogens in commercial chow: Implications for pathways influenced by hormones. *Laboratory Investigation* 81(5): 735-747, 2001.
- Ososki AL, Kennelly EJ. Phytoestrogens: a review of the present state of research. *Phytotherapy Research* 17: 845-869, 2003.
- Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT). www.food.gov.uk/multimedia/pdfs/phytoestrogenreport0503, 2003.
- Jensen MN and Ritskes-Hoitinga M. How isoflavone levels in common rodent diets can interfere with the value of animal models and with experimental results. *Laboratory Animals* 41: 1-18, 2007.
- Thigpen JE, Setchell KD, Saunders HE, Haseman JK, Grant MG, Forsy DB. Selecting the appropriate rodent diet for endocrine disruptor research and testing studies. *ILAR Journal* 45(4): 401-416, 2004.
- Yildiz F. *Phytoestrogens in Functional Foods*. CRC Press, Inc., 2006.
- Barnes S. Effects of genistein on in vitro and in vivo models of cancer. *Journal of Nutrition* 125: 777S-783S, 1995.
- Barnes S, Grubbs C, Setchell KDR, and Carlson J. Soybeans inhibit mammary tumors in models of breast cancer. *Prog. Clinical Biology Research* 347: 239-253, 1990.
- Barnes S, Peterson TG, Grubbs C, and Setchell KDR. Potential role of dietary isoflavones in the prevention of cancer. *Advances in Experimental Medicine and Biology* 354: 135-147, 1994.
- Fitzsimmons JTR, Orson NV, and El-Aaaser. Effects of soybeans and ascorbic acid on experimental carcinogenesis. *Comparative Biochemistry and Physiology* 93A: 285-290, 1989.
- Cho TM, Peng, N, Clark JT, Novak L, Roysommuti S, Prasain J, and Wyss JM. Genistein attenuates the hypertensive effects of dietary NaCl in hypertensive male rats. *Endocrinology* 148(11): 5396-5402, 2007.
- Bakhit RM, Klein BP, Essex-Sorlie D, Ham JO, Erdman JW, and Potter SM. Intake of 25 g of soybean protein with or without soybean fiber alters plasma lipids in men with elevated cholesterol concentrations. *Journal of Nutrition* 124: 213-222, 1994.
- Carroll KK and Kurowska EM. Soy consumption and cholesterol reduction: review of animal and human studies. *Journal of Nutrition* 125: 594S-597S, 1995.
- Ascencio C, Torres N, Isoard-Acosta E, Gomez-Perez FJ, Hernandez-Pando R, and Tovar AR. Soy affects serum insulin and hepatic SREBP-1 mRNA and reduces fatty liver in rats. *Journal of Nutrition* 134:522-529, 2004.
- Torre-Villalvazo I, Tovar AR, Ramos-Barragan VE, Cerbon-Cervantes MA, and Torres N. Soy protein ameliorates metabolic abnormalities in liver and adipose tissue of rats fed a high fat diet. *Journal of Nutrition* 138: 462-468, 2008.
- Cederroth CR, Vinciguerra M, Gjinovci A, Kuhne F, Klein M, Cederroth M, Caille D, Suter M, Newmann D, James RW, Doerge DR, Wallimann T, Meda P, Foti M, Rohner-Jeanrenaud F, Vassalli J, and Nef S. Dietary phytoestrogens activate AMP-activated protein kinase with improvement in lipid and glucose metabolism. *Diabetes* 57: 1176-1185, 2008.
- Messina M, Ho S, Alekel DL. Skeletal benefits of soy isoflavones: A review of the clinical trial and epidemiological data. *Current Opinion in Clinical Nutrition and Metabolic Care* 7: 649-658, 2004.
- Picherit C, Chanteranne B, Bennetau-Pelissero C, Davicco MJ, Lebecque P, Barlet JB, and Coxam V. Dose-dependent bone-sparing effects of dietary isoflavones in the ovariectomized rat. *British Journal of Nutrition* 85(3): 307-316, 2001.
- Casanova M, You L, Gaido KW, Archibeque-Engle, Janszen DB, and Heck, H. Developmental effects of dietary phytoestrogens in Sprague-Dawley rats and interactions of genistein and daidzein with rat estrogen receptors alpha and beta in vitro. *Toxicological Sciences* 51: 236-244, 1999.
- Weber KS, Setchell KDR, Stocco DM, and Lephart ED. Dietary soy-phytoestrogens decrease testosterone levels and prostate weight without altering LH, prostate 5 alpha-reductase or testicular steroidogenic acute regulatory peptide levels in adult male Sprague-Dawley rats. *Journal of Endocrinology* 170: 591-599, 2001.
- Wisniewski AB, Klein SL, Lakshmanan Y, and Gearhart JP. Exposure to genistein during gestation and lactation demasculinizes the reproductive system in rats. *Journal of Urology* 169: 1582-1586, 2003.
- Lephart ED, Setchell KD, Handa RJ, and Lund TD. Behavioral effects of endocrine-disrupting substances: phytoestrogens. *ILAR Journal* 45:443-454, 2004.
- Johnston I and Williamson G. *Phytochemical Functional Foods*. CRC Press, Inc., 2003.